

## ORIGINAL ARTICLE

# Long-term adalimumab efficacy in patients with moderate-to-severe hidradenitis suppurativa/acne inversa: 3-year results of a phase 3 open-label extension study

Christos C. Zouboulis, MD, PhD,<sup>a</sup> Martin M. Okun, MD,<sup>b</sup> Errol P. Prens, MD, PhD,<sup>c</sup> Robert Gniadecki, MD, PhD,<sup>d</sup> Peter A. Foley, MBBS, BMedSc, MD, FACD,<sup>e</sup> Charles Lynde, MD, FRCPC,<sup>f</sup> Jamie Weisman, MD,<sup>g</sup> Yihua Gu, MS,<sup>h</sup> David A. Williams, MD, MPH,<sup>h</sup> and Gregor B. E. Jemec, MD, DMSc<sup>i</sup>  
*Dessau, Germany; Fort Atkinson, Wisconsin; Rotterdam, Netherlands; Copenhagen, Denmark; Melbourne, Australia; Markham, Ontario, Canada; Atlanta, Georgia; North Chicago, Illinois; Roskilde, Denmark*

From the Departments of Dermatology, Venereology, Allergology, and Immunology, Dessau Medical Center, Brandenburg Medical School Theodor Fontane, Dessau<sup>a</sup>; Fort HealthCare, Fort Atkinson<sup>b</sup>; Department of Dermatology, Erasmus University Medical Centre Rotterdam, Rotterdam<sup>c</sup>; Department of Dermatology, Bispebjerg Hospital, University of Copenhagen<sup>d</sup>; Department of Medicine (Dermatology), The University of Melbourne, St Vincent's Hospital Melbourne, and Probit Medical Research, Skin & Cancer Foundation Inc, Melbourne<sup>e</sup>; The Lynde Centre for Dermatology and Probit Medical Research, Markham, Ontario<sup>f</sup>; Advanced Medical Research, PC, Atlanta<sup>g</sup>; AbbVie Inc, North Chicago<sup>h</sup>; and Department of Dermatology, Zealand University Hospital, Health Sciences Faculty, University of Copenhagen, Roskilde, Denmark.<sup>i</sup>

Funding sources: Supported by AbbVie, Inc.

Disclosure: Dr Zouboulis has received honoraria from AbbVie Inc, Bayer Healthcare, Biogen, and PPM for his participation as an advisor and speaker; he has also received honoraria from Allergan, Almirall, Celgene, GlaxoSmithKline, Inflarx, Novartis, and UCB for his participation as an advisor and from Jenapharm and Pierre Fabre as a speaker. His department has received grants from AbbVie Inc, AOTI, AstraZeneca, Biogen, Celgene, Dr. Reddy's, Galderma, Novartis, and UCB for his participation as an investigator. Dr Okun has received honoraria from AbbVie Inc for advisory board participation, speaker services, and consultant services, as well as from Crescendo Biosciences and Gilead for consultant services; he was an AbbVie employee at the initiation of this study. Dr Prens has received honoraria from AbbVie Inc, Amgen, AstraZeneca, Biogen, Celgene, Janssen, Leo Pharma, Novartis, Pfizer, and UCB as an advisor and speaker; his department has received investigator-initiator grants from AstraZeneca, Celgene, Janssen, and Pfizer. Dr Gniadecki has received honoraria from AbbVie Inc, Janssen, Novartis, and Amgen for participation on advisory boards and for services as an investigator and a speaker; his department has received grants from AbbVie Inc, Janssen, and Novartis for his participation as an investigator. Dr Foley has served as a consultant, investigator, speaker, and/or advisor for and/or received travel grants from Galderma, LEO/Peplin, Ascent, Clinuvel, Cutanea, Celtaxsys, Dermira, Janssen-Cilag, Eli Lilly and Company, Australian Ultraviolet Services, Roche, CSL, 3M/iNova/Valeant, GSK/Stiefel, Abbott/AbbVie, Biogen Idec, Merck Serono, Schering-Plough/MSD, Wyeth/Pfizer, Amgen,

Novartis, Celgene, Aspen, Boehringer Ingelheim, Regeneron, Sanofi Genzyme, Sun Pharma, UCB, and Bristol-Myers Squibb. Dr Lynde has received honoraria as a principal investigator, speaker, and consultant for AbbVie Inc, Amgen, Boehringer Ingelheim, Celgene, Eli Lilly and Company, Janssen, Leo Pharma, Merck, Novartis, and Regeneron. Dr Weisman has received research grants for investigator services from AbbVie Inc, Allergan, Amgen, Astra Zeneca, Boehringer Ingelheim, Braintree, Celgene, Eli Lilly and Company, Glaxo Smith Klein, Janssen, Leo Pharma, Merck, Novartis, Pfizer, Regeneron, Steifel, and Tigercat; he has also received honoraria for service on advisory boards and speaker's bureaus from AbbVie Inc, Amgen, Celgene, Eli Lilly and Company, and Janssen. Ms Gu and Dr Williams are employees of AbbVie Inc and may hold stock and/or stock options. Dr Jemec has acted as investigator for AbbVie, Actelion, Coloplast, Janssen-Cilag, Pfizer, Pierre Fabre, Inflarx, Merck Sharp & Dome, Galderma, Leo Pharma, Novartis, Regeneron, and UCB, and he has received honoraria for speaking and/or advisory boards from AbbVie, Coloplast, Pierre Fabre, Inflarx, Merck Sharp & Dome, Galderma, Leo Pharma, Novartis, and UCB, and research grants from AbbVie, Leo Pharma, and Novartis.

In addition to funding this study, AbbVie, Inc, participated in the study design; study research; collection, analysis, and interpretation of data; and writing, review, and approval of this article for publication. All authors had access to the data and participated in the development, review, and approval of the article, as well as in the decision to submit it for publication.

Some of the data reported in this article were presented at the American Academy of Dermatology Annual Meeting in Orlando, Florida, March 3-7, 2017.

Accepted for publication May 21, 2018.

Reprints not available from the authors.

Correspondence to: Christos C. Zouboulis, MD, PhD, Departments of Dermatology, Venereology, Allergology, and Immunology, Dessau Medical Center, Brandenburg Medical School Theodor Fontane, Auenweg 38, 06847 Dessau, Germany. E-mail: [christos.zouboulis@klinikum-dessau.de](mailto:christos.zouboulis@klinikum-dessau.de).

Published online July 24, 2018.

0190-9622/\$36.00

© 2018 by the American Academy of Dermatology, Inc.

<https://doi.org/10.1016/j.jaad.2018.05.040>

**Background:** The optimal long-term dosing strategy for adalimumab (ADA) in hidradenitis suppurativa/acne inversa (HS) was evaluated by pooling the results of the PIONEER phase 3 trials and an open-label extension (OLE) study.

**Objective:** To assess the response to and tolerability of long-term administration of ADA in HS.

**Methods:** The durations of the PIONEER I/II periods A, B, and OLE were 12, 24, and 52 or more weeks, respectively. Patients who entered the OLE and received ADA (40 mg every week continuously) and responders plus partial responders (PRRs) were evaluated. Primary efficacy assessments included measurement of HS clinical response (HiSCR), lesion counts, skin pain, and Dermatology Life Quality Index (DLQI). Treatment-emergent adverse events were assessed.

**Results:** At week 12, 52.3% of those receiving ADA weekly and 73.0% of PRRs achieved HiSCR. Achievement of HiSCR was maintained through week 168 in 52.3% of patients who received ADA weekly and 57.1% of PRRs. Sustained improvement in lesion counts, skin pain, and DLQI score were also observed. The safety profile throughout the OLE was similar to the profiles observed in the PIONEER studies.

**Limitations:** The OLE was uncontrolled.

**Conclusion:** Continuous weekly dosing with ADA, 40 mg, is a reasonable treatment option for long-term control of moderate-to-severe HS. (J Am Acad Dermatol <https://doi.org/10.1016/j.jaad.2018.05.040>.)

**Key Words:** acne inversa; adalimumab; continuous; dose; hidradenitis suppurativa; longitudinal; long-term; weekly.

Hidradenitis suppurativa/acne inversa (HS) is a chronic, inflammatory, recurrent debilitating skin disease of the terminal hair follicle that usually occurs after puberty and presents as painful, deep-seated, inflamed lesions in the apocrine gland-bearing areas of the body, most commonly, the axillae, inguinal, and anogenital regions.<sup>1</sup> Approximately 0.1% to 1% of the general populations in North America and Europe are affected by HS.<sup>2</sup>

HS lesions include inflammatory nodules and abscesses, tunnel (fistula and sinus) formation with purulent drainage, and subsequent scarring, all of which can affect patient health and quality of life.<sup>2-8</sup>

Adalimumab (ADA) is a monoclonal antibody against tumor necrosis factor- $\alpha$  that has been approved in Europe, Australia, and North America to treat moderate-to-severe HS, making it the only currently approved medication for this indication.<sup>9-11</sup> In 2 pivotal phase 3 randomized controlled trials, PIONEER I and II, ADA, 40 mg weekly, resulted in significantly higher clinical response rates than with placebo in patients with moderate-to-severe HS.<sup>12</sup>

## CAPSULE SUMMARY

- Adalimumab is approved for moderate-to-severe hidradenitis suppurativa/acne inversa.
- A sustained response is seen through week 168 in 52.3% of patients treated with adalimumab, 40 mg weekly, with no additional safety issues identified.
- Adalimumab can be considered for long-term control of moderate-to-severe hidradenitis suppurativa.

The PIONEER I and II trials were similarly designed phase 3, multicenter trials with 2 double-blind, placebo-controlled periods.<sup>12</sup> A phase 3, open-label extension (OLE) trial (NCT01635764) designed to assess the long-term tolerability of and response to ADA in patients with moderate-to-severe HS followed both PIONEER trials. This analysis reports the long-term results for patients who received ADA, 40 mg weekly, throughout PIONEER I and II

(36 weeks) and onward through the OLE trial to at least week 60.

## METHODS

### Patients

The PIONEER I and II patient eligibility criteria are described in detail elsewhere.<sup>12</sup> Briefly, patients were eligible if they were at least 18 years of age, had not previously received anti-tumor necrosis factor- $\alpha$  treatment, and had moderate-to-severe HS (total abscess and inflammatory nodule [AN] count of  $\geq 3$  at baseline and HS lesions in 2 distinct body areas, 1 of which was classified as Hurley stage II or III) with inadequate response to oral antibiotic treatment. In PIONEER I, patients receiving oral

*Abbreviations used:*

|        |  |
|--------|--|
| ADA:   | adalimumab                                 |
| AN:    | abscess and inflammatory nodule            |
| DLQI:  | Dermatology Life Quality Index             |
| HiSCR: | hidradenitis suppurativa clinical response |
| HS:    | hidradenitis suppurativa                   |
| OLE:   | open-label extension                       |
| PRRs:  | responders plus partial responders         |

antibiotic agents for HS were required to discontinue treatment for at least 28 days before the baseline visit. In PIONEER II, patients could continue treatment with antibiotics (doxycycline or minocycline only) in stable doses.

To be eligible for the OLE trial, patients must have participated in and completed periods A and B of PIONEER I or II. Two populations were evaluated in this study: (1) the population that received ADA treatment weekly, which included all patients who received continuous weekly ADA treatment in PIONEER I or PIONEER II and the OLE trial, and (2) the population of responders plus partial responders (PRRs), a subset of patients in the population of those receiving ADA weekly who either achieved HiSCR at week 12 (responders) or did not achieve HiSCR but achieved at least a 25% reduction in AN count relative to baseline (partial responders) at the end of period A in PIONEER I or II.<sup>13</sup>

### Study design and treatment

PIONEER I and PIONEER II and the OLE trial were conducted in accordance with Good Clinical Practice Guidelines as defined by the International Conference on Harmonization, the Declaration of Helsinki, and/or all applicable federal and local regulations.<sup>12</sup> The study protocols were approved by the independent ethics committee or institutional review board at each study site. All patients provided written informed consent before enrollment.

The study designs for PIONEER I and II are described in detail elsewhere (Fig 1).<sup>12,13</sup> In period A, patients were randomized to receive ADA, 40 mg weekly, or placebo for 12 weeks. In period B, patients who were randomized to receive ADA in period A were reassigned to receive ADA, 40 mg weekly or every other week, or placebo for 24 weeks. Patients who were randomized to receive placebo in period A were reassigned to continue receiving placebo (PIONEER II) or receive ADA, 40 mg weekly, in period B (PIONEER I).<sup>12</sup> In the OLE trial, all patients received ADA, 40 mg weekly, for at least 60 weeks or until marketing authorization or

permanent withdrawal of the marketing application in the patient's country of residence.

### Assessments


The primary objective of this study was to assess the long-term safety, tolerability, and response to continuous weekly ADA in patients with moderate-to-severe HS.

**Efficacy assessment.** Several efficacy variables were measured at specified weeks throughout the OLE study. Achievement of a hidradenitis suppurativa clinical response (HiSCR), which was defined as at least a 50% reduction from baseline in the total AN count with no increase in abscess or draining fistula count relative to baseline of the prior phase 3 study (determined at each visit), was assessed from week 2 through week 168.<sup>14</sup> Mean improvements from baseline in abscess, fistula (draining and nondraining), nodule (inflammatory and noninflammatory), and AN counts were also determined for each patient from week 2 through week 168, as was the incidence of flare, which was defined as a 25% or greater increase in the total AN count with an increase of at least 2 relative to baseline. The Patient Global Assessment of Skin Pain (numeric rating scale of 0 to 10) and modified Sartorius score<sup>15</sup> (with higher scores [no upper limit] indicating increasingly severe disease) were assessed from week 2 through week 168. Mean improvement in Dermatology Life Quality Index (DLQI) score and achievement of a DLQI score of 0 or 1 were assessed from week 4 through week 72.

**Safety assessment.** Safety assessments included recording of treatment-emergent adverse events, clinical laboratory tests, physical examinations, and vital sign measurements. All treatment-emergent adverse events were summarized by using the Medical Dictionary for Regulatory Activities (version 19.0).

### Statistical analysis

The PRR population was defined post hoc to identify the most clinically appropriate population for continuous weekly ADA dosing in period B. Multiple predictive models were applied to search for the subpopulation of interest by evaluating candidate variables, including stratification factors, patient initial response at the end of period A, and key demographic variables. The final PRR subpopulation was proposed on the basis of a threshold-based model,<sup>16,17</sup> which showed highly stable performance through a rigorous cross-validation framework.<sup>13</sup> Other predictive models were used as sensitivity analyses and



**Fig 1.** Hidradenitis suppurativa. Study design for PIONEER I, PIONEER II, and the open-label extension. <sup>a</sup>Stratified by baseline Hurley Stage II versus III (PIONEER I and II) and baseline concomitant antibiotic use (PIONEER II). <sup>b</sup>Dosing strategy: 160 mg at week 0; 80 mg at week 2; 40 mg from week 4. <sup>c</sup>Re-randomization at entry to period B, stratified by week 12 hidradenitis suppurativa clinical response (HiSCR) status and baseline Hurley stage II versus III. <sup>d</sup>Dosing strategy: 160 mg at week 12; 80 mg at week 14; 40 mg from week 16. *ADAew*, Adalimumab every week; *PBO*, placebo; *PRO*, patient-reported outcomes; *PRRs*, responders plus partial responders (a subset of patients in the population treated with ADA weekly who either achieved HiSCR at week 12 [responders] or did not achieve HiSCR but did achieve a  $\geq 25\%$  reduction in abscess and inflammatory nodule count relative to baseline [partial responders] at the end of period A in PIONEER I or II).

identified similar subpopulations that support the validity of our final results.<sup>13</sup>

This was an OLE trial, and no formal inferential statistical tests were conducted; descriptive statistics are provided. Continuous variables were summarized in terms of the number of observations, mean, standard deviation, median, minimum, and maximum; discrete variables were summarized in terms of counts and percentages. The analysis was performed with SAS statistical software (SAS Institute Inc, Cary, NC). Missing values for categorical variables were handled by nonresponder imputation in periods A and B of PIONEER I and II. The nonresponder imputation analysis categorized any patient who had a missing value at a specific visit as a nonresponder for that visit. Last observation carried forward analyses, which used the completed evaluation from the previous visit to impute missing data at later visits, was performed. Results were reported in terms of study weeks, which consisted of PIONEER I or PIONEER II and OLE study weeks, shown consecutively.

## RESULTS

### Patients

In this analysis of results across PIONEER I and II (pooled data) into the OLE trial, 88 patients comprised the ADA weekly population and 63 comprised the PRR population. All patients were followed for at least 96 weeks (36 weeks in PIONEER I or PIONEER II and at least 60 weeks in the OLE), but follow-up time varied depending on when patients entered the OLE. Key demographics are summarized in Table I. The population treated with ADA weekly was mostly female (64%) and white (92%), with a median age of 36 years and median duration of HS of 10 years. The PRR population was similar to the population treated with ADA weekly, but with a numerically lower median C-reactive protein level. The median AN, abscess, inflammatory nodule, and draining fistula counts were 9, 1, 7, and 2, respectively, and the level of patient-reported daily pain was 4.6 in the population treated with ADA weekly and 4.7 in the PRR population.

In all, 51 patients in the population treated with ADA weekly discontinued study treatment: 22

**Table I.** Baseline characteristics of the population treated with ADA weekly and the PRR population

| Characteristic                             | ADA qwk<br>(n = 88) | PRR population<br>(n = 63) |
|--|---------------------|----------------------------|
| Sex, n (%)                                 |                     |                            |
| Female                                     | 56 (63.6)           | 41 (65.1)                  |
| Male                                       | 32 (36.4)           | 22 (34.9)                  |
| Race; n (%)                                |                     |                            |
| White                                      | 81 (92.0)           | 57 (90.5)                  |
| Black                                      | 4 (4.5)             | 3 (4.8)                    |
| Other*                                     | 3 (3.4)             | 3 (4.8)                    |
| Median age, y (min, max)                   | 35.5 (18.0, 64.0)   | 36.0 (18.0, 64.0)          |
| Median BMI, kg/m <sup>2</sup> (min, max)   | 31.4 (20.3, 54.5)   | 31.6 (20.3, 49.6)          |
| Current nicotine use, n (%)                | 52 (59.1)           | 40 (63.5)                  |
| Hurley stage, n (%)                        |                     |                            |
| II   | 42 (47.7)           | 32 (50.8)                  |
| III  | 46 (52.3)           | 31 (49.2)                  |
| Median modified Sartorius score (min, max) | 103.0 (20, 1093)    | 102.0 (27, 1093)           |
| Family history of HS, n (%)                | 27 (30.7)           | 20 (31.7)                  |
| Median duration of HS, y (min, max)        | 10.34 (1.0, 40.4)   | 10.10 (1.0, 40.4)          |
| HS lesions; median (min, max)              |                     |                            |
| AN   | 9.0 (3, 71)         | 9.0 (3, 49)                |
| Abscess                                    | 1.0 (0, 13)         | 1.0 (0, 13)                |
| Draining fistula                           | 2.0 (0, 19)         | 2.0 (0, 16)                |
| Inflammatory nodule                        | 7.0 (0, 69)         | 7.0 (0, 48)                |
| Median daily pain at worst (min 0, max 10) | 4.6 (0.0, 9.7)      | 4.7 (0.0, 9.7)             |
| Median hs-CRP level, mg/L (min, max)       | 6.50 (0.2, 189.0)   | 4.8 (0.2, 66.6)            |

ADA, Adalimumab; AN, sum of abscesses and inflammatory nodules; BMI, body mass index; HS, hidradenitis suppurativa; hs-CRP, high-sensitivity C-reactive protein; max, maximum; min, minimum; PRRs, responders plus partial responders; qwk, every week.

\*Other includes Asian, American Indian/Alaska Native, Native Hawaiian or other Pacific Islander, multirace, and other races.

primarily because of adverse events or lack of efficacy, 8 because of protocol deviations or loss to follow-up, and 21 because of withdrawn consent or for other reasons (eg, pregnancy).

### Efficacy

**HiSCR.** For the 316 patients who were randomized to receive ADA weekly in the double-blind, placebo-controlled period A of the 2 PIONEER trials, the pooled data indicated achievement of HiSCR at week 12 in 50.6% of patients (160 of 316), which was significantly higher than the rate for the 317 patients randomized to receive placebo (26.8% [85 of 317]) ( $P < .001$ ).


Among the 88 patients who entered the OLE trial and were included in the population treated with ADA weekly, the percentage achieving HiSCR during period A (52.3% at week 12) was maintained throughout period B (62.5% at week 36) and the OLE trial period (52.3% at week 168) (Fig 2). The achieved HiSCR rate during period A in the PRR population (73.0% at week 12) was reduced after week 36, but the result was generally maintained

throughout period B (68.3% at week 48) and the OLE trial period (57.1% at week 168 [Fig 2]).


### Additional end points

The AN count, draining fistula count, inflammatory nodule count, and total fistula count (sum of the draining fistulas and nondraining fistulas) improved from baseline in both populations, and the response was maintained through study week 168 (Fig 3, A-D). The rates of flare (Fig 4) and a 25% or greater increase in total fistula count (Supplemental Fig 1; available at <http://www.jaad.org>) were low and stable. In addition, there was an improvement from baseline in pain, which was indicated by a mean percent decrease in numeric rating scale scores and remained generally stable in both populations through week 168 (Fig 5).

In both populations, there was also a clinically meaningful improvement (defined as a 4-point improvement by Basra et al)<sup>18</sup> in DLQI score from baseline through week 72 (5.1-6.8 points in the population treated with ADA weekly and 6.0-7.6 points in the PRR population, respectively) (Fig 6, A). The percentage of patients who achieved




**Fig 2.** Hidradenitis suppurativa. Achievement of a hidradenitis suppurativa clinical response. <sup>a</sup>Weeks include PIONEER I or PIONEER II and the open-label extension, listed consecutively. ADAew, Adalimumab every week; LOCF, last observation carried forward; PRRs, responders plus partial responders.




**Fig 3.** Hidradenitis suppurativa. Improvement in lesion count: abscesses and inflammatory nodules (A), draining fistulas (B), total fistulas (C), and inflammatory nodules (D). <sup>a</sup>Weeks include PIONEER I or PIONEER II and the open-label extension, listed consecutively. Mean improvement is relative to baseline. ADAew, Adalimumab every week; LOCF, last observation carried forward; PRRs, responders plus partial responders.

a DLQI score of 0 or 1 increased from baseline through week 48 and was generally maintained through week 72 (Fig 6, B). Similar results were observed with modified Sartorius scores (Supplemental Fig 2; available at <http://www.jaad.org>).

**Safety assessments.** The mean patient-years of exposure were 2.21 in the population treated with ADA weekly and 2.26 in the PRR population. There were no adverse events of opportunistic infections, excluding oral candidiasis; no events of active tuberculosis, lymphoma, nonmelanoma skin



**Fig 4.** Hidradenitis suppurativa. Incidence of flare (a  $\geq 25\%$  increase in the total count of abscesses and inflammatory nodules with an increase of  $\geq 2$  relative to baseline). <sup>a</sup>Weeks include PIONEER I or PIONEER II and the open-label extension, listed consecutively. ADAew, Adalimumab every week; LOCF, last observation carried forward; PRRs, responders plus partial responders.




**Fig 5.** Hidradenitis suppurativa. Improvement in skin pain. <sup>a</sup>Weeks include PIONEER I or PIONEER II and the open-label extension, listed consecutively. Mean changes are relative to baseline. Change in numeric rating scale at worst at each visit among patients with a baseline numeric rating scale score of 3 or higher. ADAew, Adalimumab every week; LOCF, last observation carried forward; PRRs, responders plus partial responders.

cancer, malignancy, or demyelinating disorder; and no deaths. Three serious infections were reported in the population treated with ADA weekly; 1 (pneumonia) was considered probably related to the treatment. The most frequently reported treatment-emergent adverse events (a rate of  $\geq 10\%$  in either population) were hidradenitis, upper respiratory tract infection, headache, nasopharyngitis, arthralgia, influenza, diarrhea, urinary tract infection, sinusitis, bronchitis, dizziness, and nausea (Table II).

## DISCUSSION

Efficacy and safety results derived from pooling the pivotal phase 3 trials with the OLE trial confirm that ADA sustains efficacy and has an acceptable safety profile following a strategy of continuous weekly treatment. Pooling of the data from the 2 PIONEER trials was prespecified in the protocol for period A and period B results. The combined analysis was appropriate on the basis of the similarities in study design and provided a more robust assessment by combining the somewhat



**Fig 6.** Hidradenitis suppurativa. Improvement in dermatology-specific quality of life: mean improvement in Dermatology Life Quality Index from baseline (**A**) and achievement of a Dermatology Life Quality Index score of 0 or 1 (no effect of skin disease on quality of life) (**B**). <sup>a</sup>Weeks include PIONEER I or PIONEER II and the open-label extension, listed consecutively. Mean changes are relative to baseline. *ADAew*, Adalimumab every week; *LOCF*, last observation carried forward; *PRRs*, responders plus partial responders.

**Table II.** Treatment-emergent adverse events

| Adverse events, n (%)                              | ADA qwk population<br>n = 88 | PRR population<br>n = 63 |
|--|------------------------------|--------------------------|
| Any event  | 76 (86.4)                    | 55 (87.3)                |
| Leading to discontinuation of the study drug       | 13 (14.8)                    | 10 (15.9)                |
| Serious  | 12 (13.6)                    | 9 (14.3)                 |
| Infections   | 63 (71.6)                    | 45 (71.4)                |
| Serious infections                                 | 3 (3.4)*                     | 2 (3.2) <sup>†</sup>     |
| Latent tuberculosis                                | 2 (2.3)                      | 2 (3.2)                  |
| Most frequently reported TEAEs (≥10%) <sup>‡</sup> |                              |                          |
| Hidradenitis                                       | 21 (23.9)                    | 16 (25.4)                |
| Upper respiratory tract infection                  | 17 (19.3)                    | 15 (23.8)                |
| Headache   | 17 (19.3)                    | 13 (20.6)                |
| Nasopharyngitis                                    | 16 (18.2)                    | 13 (20.6)                |
| Arthralgia   | 12 (13.6)                    | 10 (15.9)                |
| Influenza  | 13 (14.8)                    | 9 (14.3)                 |
| Diarrhea   | 7 (8.0)                      | 7 (11.1)                 |
| Urinary tract infection                            | 9 (10.2)                     | 4 (6.3)                  |
| Sinusitis  | 8 (9.1)                      | 6 (9.5)                  |
| Bronchitis   | 6 (6.8)                      | 6 (9.5)                  |
| Dizziness  | 6 (6.8)                      | 6 (9.5)                  |
| Nausea   | 6 (6.8)                      | 6 (9.5)                  |

ADA, Adalimumab; PRRs, responders plus partial responders; qwk, every week; TEAE, treatment-emergent adverse event.

\*Includes pneumonia (n = 2) and cellulitis of the right leg (n = 1).

<sup>†</sup>Includes pneumonia (n = 1) and cellulitis of the right leg (n = 1).

<sup>‡</sup>In either population.

limited sample sizes in period B of each study. The data for the populations of patients with HS who received weekly ADA treatment spanning the PIONEER I and II studies and the OLE trial confirm

that weekly ADA treatment maintained long-term responses, as demonstrated by achievement of HiSCR and improvements in lesion counts, skin pain, and DLQI score. HiSCR was achieved by at

least half of the patients in both the population treated with ADA weekly and the PRR population at week 12 (the primary end point of PIONEER I and II), and the HiSCR was maintained through week 168 (OLE). Study closure was at the same time for all patients, and those who enrolled later did not have the opportunity to reach the later time points.

Improvements from baseline in skin lesions, including in AN count, draining fistulas, total fistulas, inflammatory nodules, and skin pain, were observed as early as week 2, also detected at week 12, and maintained through week 168. The natural history of untreated HS is chronic and progressive in some patients.<sup>19-21</sup> Although the data presented here are uncontrolled, the mean improvement in total fistula count suggests that some patients with HS who were treated with continuous weekly ADA may have experienced a halt in progression of HS, suggesting that ADA may be disease modifying. Additionally, a clinically meaningful improvement in DLQI score was observed through week 72.

Results from the PRR population demonstrate the effect of continuous weekly dosing because findings indicate that patients who had even a partial response at week 12 were likely to achieve HiSCR if they continued treatment with ADA. Continuous weekly dosing in patients who achieved at least a 25% reduction in AN count relative to baseline at week 12 is also supported by the outcomes of other end points, including reduced levels of skin pain and lower lesion counts. Given the current absence of alternative therapies, the burden of disease in the target population, and the observation of late achievement of some of the benefits in pain relief and improved scores on DLQI assessments (weeks 48 and 72), continuation of therapy may be considered even in patients with an initial partial response after short-term treatment.

The safety profile of long-term weekly administration of ADA therapy in this analysis was similar to that observed in PIONEER I and II, demonstrating the longitudinal safety of ADA in the treatment of patients with moderate-to-severe HS. The results were consistent with the known ADA safety profile.<sup>12-22</sup> ADA was generally well tolerated, and no new safety risks were identified in this patient population.


In conclusion, on the basis of the data from the studies that have been reported here and elsewhere,<sup>12</sup> ADA dosing of 40 mg weekly is effective and safe for long-term treatment of patients with moderate-to-severe HS.

The authors would like to thank Piyalal Karunaratne for statistical analysis support and Holly Capasso-Harris, PhD, of JB Ashtin, who, with funding from and on behalf of AbbVie, developed the first draft of this article on the basis of an author-approved outline and assisted in implementing author revisions.


## REFERENCES

1. Zouboulis CC, Desai N, Emtestam L, et al. European S1 guideline for the treatment of hidradenitis suppurativa/acne inversa. *J Eur Acad Dermatol Venerol*. 2015;29(4):619-644.
2. Gulliver W, Zouboulis CC, Prens E, et al. Evidence-based approach to the treatment of hidradenitis suppurativa/acne inversa, based on the European guidelines for hidradenitis suppurativa. *Rev Endocr Metab Disord*. 2016;17(3):343-351.
3. Lipsker D, Severac F, Freys M, et al. The ABC of hidradenitis suppurativa: a validated glossary on how to name lesions. *Dermatology*. 2016;232(2):137-142.
4. Zouboulis CC, del Marmo V, Mrowietz, et al. Hidradenitis suppurativa/acne inversa: criteria for diagnosis, severity assessment, classification and disease evaluation. *Dermatol*. 2015;231(2):184-190.
5. Zouboulis CC, Tzellos T, Kyrgidis A, et al. Development and validation of IHS4, a novel dynamic scoring system to assess hidradenitis suppurativa/acne inversa severity. *Br J Dermatol*. 2017;177:1401-1409.
6. Jemec GB, Heidenheim M, Nielsen NH. Hidradenitis suppurativa—characteristics and consequences. *Clin Exp Dermatol*. 1996;21(6):419-423.
7. von der Werth JM, Jemec GB. Morbidity in patients with hidradenitis suppurativa. *Br J Dermatol*. 2001;144(4):809-813.
8. Matusiak L, Bieniek A, Szepietowski JC. Hidradenitis suppurativa markedly decreases quality of life and professional activity [letter]. *J Am Acad Dermatol*. 2010;62(4):706-708.
9. Adalimumab [package insert]. AbbVie Inc. 2016. North Chicago, IL.
10. European Medicines Agency. Humira Adalimumab. Available at: [http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/000481/human\\_med\\_000822.jsp&mid=WC0b01ac058001d124](http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/000481/human_med_000822.jsp&mid=WC0b01ac058001d124). Accessed June 30, 2018
11. Zouboulis CC. Adalimumab for the treatment of hidradenitis suppurativa/acne inversa. *Expert Rev Clin Immunol*. 2016;12(1):1015-1026.
12. Kimball AB, Okun MM, Williams DA, et al. Two phase 3 trials of adalimumab for hidradenitis suppurativa. *N Engl J Med*. 2016;375(5):422-434.
13. Gulliver W, Okun M, Martorell A, et al. Therapeutic response guided dosing strategy to optimize long-term adalimumab treatment in patients with hidradenitis suppurativa: integrated results from the PIONEER phase 3 trials [abstract]. *J Am Acad Dermatol*. 2017;76(suppl 1):5098.
14. Kimball AB, Jemec GBE, Yang M, et al. Assessing the validity, responsiveness, and meaningfulness of the hidradenitis suppurativa clinical response (HiSCR) as the clinical endpoint for hidradenitis suppurativa treatment. *Br J Dermatol*. 2014;171(6):1434-1442.
15. Sartorius K, Lapins J, Emtestam L, Jemec GE. Suggestions for uniform outcome variables when reporting treatment effects in hidradenitis suppurativa. *Br J Dermatol*. 2003;149(1):211-213.
16. Chen G, Zhong H, Belousov A, Devanarayan V. A PRIM approach to predictive-signature development for patient stratification. *Stat Med*. 2015;34(2):317-342.
17. McKeegan EM, Ansell PJ, Davis G, et al. Plasma biomarker signature associated with improved survival in advanced

- non-small cell lung cancer patients on linifanib. *Lung Cancer*. 2015;90(2):296-301.
18. Basra MK, Salek MS, Camilleri L, Sturkey R, Finlay AY. Determining the minimal clinically important difference and responsiveness of the Dermatology Life Quality Index (DLQI): further data. *Dermatology*. 2015;230(1):27-33.
  19. Micheletti RG. Natural history, presentation, and diagnosis of hidradenitis suppurativa. *Semin Cutan Med Surg*. 2014; 33(suppl 3):S51-S53.
  20. Kromann CB, Deckers IE, Esmann S, Boer J, Prens EP, Jemec GB. Risk factors, clinical course and long-term prognosis in hidradenitis suppurativa: a cross-sectional study. *Br J Dermatol*. 2014;171(4):819-824.
  21. von der Werth JM, Williams JC. The natural history of hidradenitis suppurativa. *J Eur Acad Dermatol Venereol*. 2000; 14(5):389-392.
  22. Burmester GR, Panaccione R, Gordon KB, et al. Adalimumab: long-term safety in 23 458 patients from global clinical trials in rheumatoid arthritis, juvenile idiopathic arthritis, ankylosing spondylitis, psoriatic arthritis, psoriasis, and Crohn's disease. *Ann Rheum Dis*. 2013;72(4): 517-524.



**Supplemental Fig 1.** Hidradenitis suppurativa. Incidence of  $\geq 25\%$  increase in total fistula count. <sup>a</sup>Weeks include PIONEER I or II and the open-label extension, listed consecutively. Mean improvement is relative to baseline. *ADAew*, Adalimumab every week; *LOCF*, last observation carried forward; *N*, number of patients in the population; *PPR*, responders plus partial responders.



**Supplemental Fig 2.** Hidradenitis suppurativa. Mean improvement from baseline in modified Sartorius scores. <sup>a</sup>Weeks include PIONEER I or II and the open-label extension, listed consecutively. Mean improvement is relative to baseline. *ADAew*, Adalimumab every week; *LOCF*, last observation carried forward; *N*, number of patients in the population; *PPR*, responders plus partial responders.